



AP Calculus AB FRQ Practice · Unit 2

Derivatives From a Table · numberbender.com

Name: _____

Date: _____

Score: / 9

What this question practices

- Estimating a derivative with a difference quotient
- The product rule
- The quotient rule
- Recognizing the limit definition of the derivative

No calculator is allowed for this question. Show all your work, and justify your answers. Suggested time: 15 minutes.

The functions f and g are differentiable for all real numbers. The table gives selected values of f , g and their derivatives.

x	$f(x)$	$f'(x)$	$g(x)$	$g'(x)$
1	4	2	3	-1
2	7	4	5	2
3	12	6	2	-3

(a) Use the data in the table to estimate $f'(2.5)$. Show the computation that leads to your answer.

(b) Let h be the function defined below. Find $h'(2)$.

$$h(x) = f(x) g(x)$$

(c) Let k be the function defined below. Find $k'(3)$.

$$k(x) = \frac{f(x)}{g(x)}$$



(d) The limit below is the derivative of a function p at a point. Identify p and the point, and find the value of the limit.

$$\lim_{h \rightarrow 0} \frac{(2+h)^4 - 16}{h}$$





This question is worth 9 points, scored the way AP Calculus readers score free-response questions. Compare your work with the model solution, then award yourself points using the scoring notes.

(a) Use the data in the table to estimate $f'(2.5)$. Show the computation that leads to your answer.

$$f'(2.5) \approx \frac{f(3) - f(2)}{3 - 2} = \frac{12 - 7}{1} = 5$$

→ Use the average rate of change of f over the interval $[2, 3]$, which contains 2.5.

Scoring. 1 point for a difference quotient using $x = 2$ and $x = 3$, with the value 5.

(b) Let h be the function defined below. Find $h'(2)$.

$$h(x) = f(x)g(x)$$

$$h'(2) = f'(2)g(2) + f(2)g'(2) = 4(5) + 7(2) = 34$$

→ Apply the product rule, then read the values at $x = 2$ from the table.

Scoring. 2 points: 1 point for the product rule, 1 point for the answer 34.

(c) Let k be the function defined below. Find $k'(3)$.

$$k(x) = \frac{f(x)}{g(x)}$$

$$k'(3) = \frac{f'(3)g(3) - f(3)g'(3)}{(g(3))^2} = \frac{6(2) - 12(-3)}{2^2} = \frac{48}{4} = 12$$

→ Apply the quotient rule, being careful with the subtraction order and the sign of $g'(3)$.

Scoring. 3 points: 1 point for the quotient rule, 1 point for correct substitution, 1 point for the answer 12.

(d) The limit below is the derivative of a function p at a point. Identify p and the point, and find the value of the limit.

$$\lim_{h \rightarrow 0} \frac{(2+h)^4 - 16}{h}$$

$$p(x) = x^4, \quad x = 2: \quad p'(x) = 4x^3, \quad p'(2) = 4(8) = 32$$

→ Compare with the definition of the derivative: the limit of $(p(2+h) - p(2)) / h$ as h approaches 0, where $p(2) = 16$.

Scoring. 3 points: 1 point for $p(x) = x^4$, 1 point for the point $x = 2$, 1 point for the value 32.

